

An algorithm for constructing Hjelmslev planes

Joanne L. Hall^{*} and Asha Rao[†]

^{*†} School of Mathematical and Geospatial Sciences

RMIT University

^{*}School of Mathematical Sciences

Queensland University of Technology

February 20, 2013

Abstract: Projective Hjelmslev planes and Affine Hjelmselv planes are generalisations of projective planes and affine planes. We present an algorithm for constructing a projective Hjelmslev planes and affine Hjelmselv planes using projective planes, affine planes and orthogonal arrays. We show that all 2-uniform projective Hjelmslev planes, and all 2-uniform affine Hjelmselv planes can be constructed in this way. As a corollary it is shown that all 2-uniform Affine Hjelmselv planes are sub-geometries of projective Hjelmselv planes.

1 Introduction

Hjelmslev planes are named after Danish geometer J. Hjelmslev and were first formally described by Klingenberg in the 1950s [16]. Some work was done in the 60s and 70s describing the properties of Hjelmslev planes, see for example [6, 8, 4].

A connection with linear codes [12] and a conjectured connection to mutually unbiased bases [17] has recently rekindled interest in Hjelmslev planes. Explicit

constructions and concrete examples are required for further investigation in these application areas. Hjelmslev planes can be constructed using chain rings [19]. Galois ring Hjelmslev geometries are the most thoroughly investigated of the chain ring geometries [12, 14], more recent work has included other chain rings [10].

We show an algorithm for constructing projective Hjelmslev planes, some of which cannot be calculated using rings. This algorithm is easily implemented using any computing language so that a Hjelmslev plane may be generated for use in applications. The construction uses a projective plane, an affine plane and an orthogonal array. There are open online lookup tables (eg. [1, 18]) for these objects, or some may be constructed using a Galois field [3, §VII.2]. Drake and Shult show that all Hjelmslev planes can be constructed from a projective plane and semi-nets with zings [9, Prop 2.4], however there is no library of semi-nets with zings. Algorithm 1 for constructing Projective Hjelmslev planes is implied by the proof of [13, Thm 1], but is given here in algorithmic language. Algorithm 2 for constructing affine Hjelmslev planes is new.

Section 2 contains definitions and facts about projective Hjelmslev planes. Section 3 describes the mathematical objects required for the construction. Sections 4 and 5 detail algorithms for constructing 2-uniform projective Hjelmslev planes and 2-uniform affine Hjelmslev planes. Section 6 discusses some properties of the algorithm and concludes with some further directions.

2 Definitions

Hjelmslev planes are generalisations of projective and affine planes. We write $P \in h$, to show that the point P is incident with line h ; PR is a line incident with points P and R ; $f \cap h$ is a point of intersection of lines f and h . In Hjelmslev geometry, lines (or points) may have the relationship of neighbour, denoted $g \sim h$ (or $P \sim Q$).

Definition 1. [8] A projective Hjelmslev plane, $\mathcal{H}$, is an incidence structure such that:

1. any two points are incident with at least one line.
2. any two lines intersect in at least one point.
3. any two lines g, h that meet at more than one point are neighbours, denoted $g \sim h$.
4. any two points P and Q that are incident with more than one line are neighbours, denoted $P \sim Q$.
5. there exists an epimorphism ϕ from $\mathcal{H}$ to an ordinary projective plane $\mathcal{P}$ such that:

$$(a) \quad \phi(P) = \phi(Q) \iff P \sim Q$$

$$(b) \quad \phi(g) = \phi(h) \iff g \sim h$$

The neighbour property is an equivalence relation so the set of lines is partitioned into line-neighbourhoods and the set of points into point-neighbourhoods. A projective Hjelmslev plane is denoted by $(t, r)PH$ -plane where t is the number of neighbouring points on each line, and r is the order of the projective plane associated by the epimorphism ϕ . Furthermore it is known that $r = s/t$ where $t + s$ is the number of points on each line, each line-neighbourhood has t^2 lines and each point-neighbourhood has t^2 points [8, Defi 1.5],[15]. Just as in projective planes, points and lines are dual. A $(1, r)PH$ -plane is a projective plane of order r (See Definition 5). This notation should not be confused with $PH(R)$, the projective Hjelmslev plane over the ring R , or $PH(n, p^r)$ where $R = GR(p^n, r)$ [17, 12].

Definition 2. [8] An Affine Hjelmslev plane $\mathcal{H}$ is an incidence structure such that

1. any two points are incident with at least one line.
2. any two lines g, h that meet at more than one point are neighbours, denote $g \sim h$.
3. There exists an epimorphism ϕ from $\mathcal{H}$ to an ordinary affine plane $\mathcal{A}$ such that
 - (a) $\phi(P) = \phi(Q) \iff P \sim Q$,
 - (b) $\phi(g) = \phi(h) \iff g \sim h$,
 - (c) $|g \cap h| = 0 \Rightarrow \phi(g) \parallel \phi(h)$.

An Affine Hjelmslev plane may be denoted $(t, r)AH$ -plane, where t is the number of neighbouring points on each line, and $r = s/t$ where s is the number of points on each line. An Affine Hjelmslev plane may have parallel lines which are neighbours as condition 3c is a one way implication.

Hjelmslev planes are mentioned in some books on finite geometry [5, §7.2], but not in the more standard works on design theory eg [3, 2].

Hjelmslev planes have a rich structure with several interesting substructures. The following functions allows interrogation of the structure of each point neighbourhood.

Definition 3. [6, Defi 2.3] Let P be a point of a Hjelmslev plane $\mathcal{H}$. The point-neighbourhood restriction, $\bar{P}$, is defined as follows:

1. the points of $\bar{P}$ are the points Q of $\mathcal{H}$ such that $Q \sim P$.
2. the lines of $\bar{P}$ are the restrictions of lines g of $\mathcal{H}$ to the points in $\bar{P}$:
 $g_P = g \cap \bar{P}$.

Definition 4. [6, Defi 2.4] A 1-uniform projective (affine) Hjelmslev plane $\mathcal{H}$ is an ordinary projective (affine) plane. A projective (affine) Hjelmslev plane is n -uniform if

1. for every point $P \in \mathcal{H}$, $\bar{P}$ is an $(n - 1)$ uniform affine Hjelmslev plane.
2. for each point P of $\mathcal{H}$, every line of $\bar{P}$ is the restriction of the same number of lines of $\mathcal{H}$.

In a 2-uniform projective Hjelmslev plane every point-neighbourhood restriction is an ordinary affine plane. It is known that a projective Hjelmslev plane is 2-uniform if and only if it is a (t, t) -PH plane, similarly an affine Hjelmslev plane is 2-uniform if and only if it is a (t, t) -AH plane [15]. All (t, t) -PH planes that can be generated by rings have been catalogued [12]. However there are many more Hjelmslev planes that cannot be generated by rings.

3 The objects required for the construction

We present an algorithm for generating a 2-uniform projective Hjelmslev plane. This algorithm is inspired by a construction by Drake and Schult [9, 2.4] which uses a projective plane and a semi-net with zings. Simi-nets have not been well catalogued, making Drake and Schults construction difficult to implement. The algorithms given in section 4 and 5 are restricted to (t, t) -PH planes and (t, t) -AH planes, but use objects that are well studied, and catalogued. The algorithm takes three different classes of combinatorial structures and use them to generate (t, t) -PH plane and (t, t) -AH planes. Examples and properties of the combinatorial structures can be found in any standard work on combinatorial designs eg. [3, 2].

Definition 5. [3, VII.2.2] A *projective plane* is a set of points and lines such that

- any two distinct points are incident with exactly one line.
- any two distinct lines meet at exactly one point.
- there exist four points no three of which are on a common line.

A projective plane of order m has $m + 1$ points on each line, $m + 1$ lines through each point, $m^2 + m + 1$ points and $m^2 + m + 1$ lines. A projective plane of order m may be represented as a $2 - (m^2 + m + 1, m + 1, 1)$ block design [2, Defi 2.9].

Definition 6. [3, VII.2.11] A *finite affine plane* is a set of points and lines such that

- any two distinct points are incident with exactly one line.
- for any point P not on line l there is exactly one line through P that is parallel (has no points in common) with l .
- there exist three points not on a common line.

An affine plane of order m has m points on a line, $m + 1$ lines through each point, m^2 points and $m^2 + m$ lines. An affine plane can be obtained from a projective plane by removing one line and all the points on that line. An affine plane of order m may be represented as a $2 - (m^2, m, 1)$ block design [2, Defi 2.9]. An affine plane of order m has a parallelism, each $\parallel$ -class contains m lines.

$\parallel$ -classes S and T are orthogonal if each line of S has exactly one element in common with each line of T . An affine plane of order m has $m + 1$ mutually orthogonal $\parallel$ -classes.

Definition 7. [3, III.3.5] An *orthogonal array* $OA(t, k, v)$ is a $v^2 \times k$ array with entries from a set of v symbols, such that in any t columns each ordered t -tuple occurs exactly once.

Each symbol occurs in each column of the orthogonal array v times. An orthogonal array may be obtained from an affine plane by assigning each point to a row, and each $\parallel$ -class to a column of the array. The symbol in position i, j of the array indicates the line of $\parallel$ -class j that is incident with point i .

The three structures above may all be generated from a projective plane. However, it is not essential that the objects have any relationship other than

size. There are open online lookup tables (eg. [1, 18]) for these objects, as well as many software tools for generating orthogonal arrays. Some of these objects may also be constructed using Galois fields [3, §VII.2].

The construction of Drake and Shult [9] uses the incidence matrix of a projective plane, for which each symbol is replaced by a submatrix which meets a set of conditions given by a semi-net with zings. The algorithm developed here is related to this method.

4 An algorithm for constructing 2–uniform projective Hjelmslev planes

Algorithm 1. To create the structure $\mathcal{H}$:

Step 1 Let $\mathcal{P}$ be a projective plane of order m , Let $\mathcal{A}_0, \mathcal{A}_1, \dots, \mathcal{A}_{m^2+m}$ be a list of an affine planes of order m and $\mathcal{O}_0, \mathcal{O}_1, \dots, \mathcal{O}_{m^2+m}$ be a list of an orthogonal array $OA(2, m+1, m)$. Note that some (or all) of the affine planes and orthogonal arrays may be the same.

Step 2 For each point of $\mathcal{P}$, replace point P with m^2 points which are a copy of $\mathcal{A}_P$. This gives $(m^2 + m + 1)m^2$ points in $\mathcal{H}$. Each affine plane will now be called a point-neighbourhood of $\mathcal{H}$.

Step 3 Choose a line $l = \{P_0, P_2, \dots, P_m\}$, in $\mathcal{P}$, and for each point of l choose a parallel class of each of $\mathcal{A}_{P_0}, \mathcal{A}_{P_1}, \dots, \mathcal{A}_{P_m}$. Label each of the lines of the parallel class of each point-neighbourhood with the symbols from $\mathcal{O}_l$. Since each point-neighbourhood is in m lines of $\mathcal{P}$, each time a particular point P of $\mathcal{P}$ is in a chosen line, a different parallel class of $\mathcal{A}_P$ must be used. Label each column of $\mathcal{O}_l$ with a point of l .

Step 4 Each line of $\mathcal{H}$ is constructed by reading a row of $\mathcal{O}_l$ and selecting the points which correspond the lines of the the point neighbourhoods.

$$\mathcal{P} = \left\{ \begin{array}{l} \{0, 1, 2, 9\}, \\ \{3, 4, 5, 9\}, \\ \{6, 7, 8, 9\}, \\ \{0, 3, 6, A\}, \\ \{1, 4, 7, A\}, \\ \{2, 5, 8, A\}, \\ \{0, 4, 8, B\}, \\ \{1, 5, 6, B\}, \\ \{2, 3, 7, B\}, \\ \{0, 7, 5, C\}, \\ \{1, 3, 8, C\}, \\ \{2, 4, 6, C\}, \\ \{9, A, B, C\}, \end{array} \right\} \quad \mathcal{A} = \left\{ \begin{array}{l} \{R, S, T\}, \\ \{U, V, W\}, \\ \{X, Y, Z\}, \\ \{R, U, X\}, \\ \{S, V, Y\}, \\ \{T, W, Z\}, \\ \{R, V, Z\}, \\ \{S, W, X\}, \\ \{T, U, Y\}, \\ \{R, W, Y\}, \\ \{S, U, Z\}, \\ \{T, V, X\} \end{array} \right\} \quad \mathcal{O} = \begin{array}{cccc} L & L & L & L \\ L & M & M & M \\ L & N & N & N \\ M & L & M & N \\ M & M & N & L \\ M & N & L & M \\ N & L & N & M \\ N & M & L & N \\ N & N & M & L \end{array}$$

Figure 1: Step 1. A Projective plane of order 3, and Affine plane of order 3 and an orthogonal array $OA(2, 4, 3)$. In this example $\mathcal{A}_i = \mathcal{A}$ and $\mathcal{O}_i = \mathcal{O}$ for all $0 \leq i \leq m^2 + m$.

Repeat for each line of $\mathcal{P}$.

An example is given in figures 1-4.

Theorem 1. *The structure generated by Algorithm 1 is a 2-uniform (m, m) PH-plane.*

Proof. This algorithm generates an incidence structure $\mathcal{H}$ with $(m^2 + m + 1)m^2$ points, $(m^2 + m + 1)m^2$ lines, each line containing $(m^2 + m)$ points, and each point incident with $(m^2 + m)$ lines.

Axioms 1 and 4: Any pair of points P and Q which are in the same point-neighbourhood are incident with exactly one line of the point neighbourhood restriction, which is an affine plane. Each line of the point neighbourhood re-

$$\{(0, R), (0, S), (0, T), (0, U), (0, V), (0, W), (0, X), \dots, (C, X), (C, Y), (C, Z)\}.$$

Figure 2: Step 2. The points of $\mathcal{H}$ can be written as a Cartesian product of the set of points in $\mathcal{A}$ and $\mathcal{P}$

In neighbourhood 3; $L := \{R, S, T\}$, $M := \{U, V, W\}$, $N := \{X, Y, Z\}$.

In neighbourhood 4; $L := \{R, S, T\}$, $M := \{U, V, W\}$, $N := \{X, Y, Z\}$.

In neighbourhood 5; $L := \{R, S, T\}$, $M := \{U, V, W\}$, $N := \{X, Y, Z\}$.

In neighbourhood 9; $L := \{R, U, X\}$, $M := \{S, V, Y\}$, $N := \{T, W, Z\}$.

3	4	5	9
L	L	L	L
L	M	M	M
L	N	N	N
M	L	M	N
M	M	N	L
M	N	L	M
N	L	N	M
N	M	L	N
N	N	M	L

Figure 3: Step 3. Choosing line $l = \{3, 4, 5, 9\}$, the chosen $\parallel$ -classes of each point neighbourhood of l , and the labels for the columns of $\mathcal{O}$.

$$\begin{aligned}
& \{(3, R), (3, S), (3, T), \quad (4, R), (4, S), (4, T), \quad (5, R), (5, S), (5, T), \quad (9, R), (9, U), (9, X)\} \\
& \{(3, R), (3, S), (3, T), \quad (4, U), (4, V), (4, W), \quad (5, U), (5, V), (5, W), \quad (9, S), (9, V), (9, Y)\} \\
& \{(3, R), (3, S), (3, T), \quad (4, X), (4, Y), (4, Z), \quad (5, X), (5, Y), (5, Z), \quad (9, T), (9, W), (9, Z)\} \\
& \{(3, U), (3, V), (3, W), \quad (4, R), (4, S), (4, T), \quad (5, U), (5, V), (5, W), \quad (9, T), (9, W), (9, Z)\} \\
& \{(3, U), (3, V), (3, W), \quad (4, U), (4, V), (4, X), \quad (5, X), (5, Y), (5, Z), \quad (9, R), (9, U), (9, X)\} \\
& \{(3, U), (3, V), (3, W), \quad (4, X), (4, Y), (4, Z), \quad (5, R), (5, S), (5, T), \quad (9, S), (9, V), (9, Y)\} \\
& \{(3, X), (3, Y), (3, Z), \quad (4, R), (4, S), (4, T), \quad (5, X), (5, Y), (5, Z), \quad (9, S), (9, V), (9, Y)\} \\
& \{(3, X), (3, Y), (3, Z), \quad (4, U), (4, V), (4, X), \quad (5, R), (5, S), (5, T), \quad (9, T), (9, W), (9, Z)\} \\
& \{(3, X), (3, Y), (3, Z), \quad (4, X), (4, Y), (4, Z), \quad (5, U), (5, V), (5, W), \quad (9, R), (9, U), (9, X)\}
\end{aligned}$$

Figure 4: Step 4. The lines of $\mathcal{H}$ in the line-neighbourhoods corresponding to the line $\{3, 4, 5, 9\}$ and $\{6, 7, 8, 9\}$ of $\mathcal{P}$ are constructed according to $\mathcal{O}$. Note that every pair of lines from within a line neighbourhood share exactly 3 points.

striction is used in m lines of $\mathcal{H}$, as each symbol appears m times in each column of $\mathcal{O}$. For points P and R which are in different point-neighbourhoods, there is exactly one line of $\mathcal{P}$ which is incident with any pair of point-neighbourhoods. Given $\parallel$ -classes $\bar{P}_X$ and $\bar{R}_Y$ of each point-neighbourhood, $\mathcal{O}$ ensures that each line of $\bar{P}_X$ is in a line of $\mathcal{H}$ with each line of $\bar{R}_Y$ exactly once.

Axioms 2 and 3: $\mathcal{O}$ ensures that lines in the same line neighbourhood meet in exactly one line of a $\parallel$ -class of a point-neighbourhood, which is m points. For lines g and h which are in different line-neighbourhoods, their line-neighbourhoods may be labeled with lines from $\mathcal{P}$. Any pair of lines in $\mathcal{P}$ intersect in exactly one point, thus any line-neighbourhoods of $\mathcal{H}$ intersect in exactly one point neighbourhood $\bar{Q}$. Each line-neighbourhood is allocated a different $\parallel$ -class $\bar{Q}_X, \bar{Q}_Y$. Thus the line g in $\mathcal{H}$ must contain a line of $\bar{Q}_X$ and h a line of $\bar{Q}_Y$. As the $\parallel$ -classes are orthogonal, g and h meet in exactly one point.

Let ϕ collapse point-neighbourhoods and line neighbourhoods. It is trivial to

check that this is incidence preserving and surjective, and thus an epimorphism.

All axioms of Definition 1 are satisfied. Thus $\mathcal{H}$ is a projective Hjelmslev plane.

The cardinalities of $\mathcal{H}$ show that it is a (m, m) -PH-plane and is therefore 2-uniform. $\square$

In the example the affine plane used is a sub-geometry of the projective plane. However this is not required and an affine plane that is not a sub-geometry of the projective plane may be used. Any projective plane, any affine plane and any orthogonal array of the appropriate sizes may be used. In fact different affine planes may be used for each point-neighbourhood.

Theorem 2. *All 2-uniform projective Hjelmslev planes can be generated using Algorithm 1.*

Proof. The proof of Theorem 1 is easily adapted to Algorithm 2, so $\mathcal{H}$ is a 2-uniform projective Hjelmslev plane. Axiom 1 of Definition 4 requires that point-neighbourhood restrictions are affine planes. This algorithm delivers. Requiring that every line of every $\bar{P}$ is the restriction of the same number of lines is equivalent to ensuring that each line of each parallel class of the point-neighbourhood is included in the same number of lines at step 4. This is ensured as each symbol occurs in each column of an orthogonal array the same number of times. $\square$

5 An Algorithm for constructing 2-uniform affine Hjelmslev planes

Algorithm 1 may be amended to construct 2-uniform affine Hjelmslev planes

Algorithm 2. To create the structure $\mathcal{H}$:

Step 1 Let $\mathcal{A}$ be a affine plane of order m , Let $\mathcal{A}_0, \mathcal{A}_1, \dots, \mathcal{A}_{m^2+m}$ be a list of an affine planes of order m and $\mathcal{O}_0, \mathcal{O}_1, \dots, \mathcal{O}_{m^2+m}$ be a list of an orthogonal array $OA(2, m, m)$. Note that some (or all) of the affine planes and orthogonal arrays may be the same.

Step 2 For each point of $\mathcal{A}$, replace point P with m^2 points which are a copy of $\mathcal{A}_P$. This gives m^4 points in $\mathcal{H}$. Each affine plane will now be called a point-neighbourhood of $\mathcal{H}$.

Step 3 Choose a line $l = \{P_0, P_2, \dots, P_{m-1}\}$, in $\mathcal{H}$, and for each point of l choose a parallel class of each of $\mathcal{A}_{P_0}, \mathcal{A}_{P_1}, \dots, \mathcal{A}_{P_{m-1}}$. Label each of the lines of the parallel class of each point-neighbourhood with the symbols from $\mathcal{O}_l$. Since each point-neighbourhood is in m lines of $\mathcal{P}$, each time a particular point P of $\mathcal{P}$ is in a chosen line, a different parallel class of $\mathcal{A}_P$ must be used. Label each column of $\mathcal{O}_l$ with a point of l .

Step 4 Each line of $\mathcal{A}$ is constructed by reading a row of $\mathcal{O}_l$ and selecting the points which correspond the lines of the the point neighbourhoods.

Repeat for each line of $\mathcal{H}$.

Theorem 3. *All 2-uniform affine Hjelmslev planes can be generated using Algorithm 2.*

The proof is similar to that of Theorem 1 and is omitted.

Affine planes are sub-geometries of projective planes, the same is sometimes true for affine Hjelmslev planes.

Lemma 1. *[8] A $(t, r)PH$ -plane can be truncated to a $(t, r)AH$ plane*

Proof. Take $PH(t, r)$ and remove all the lines of one line-neighbourhood together with all incident points. $\square$

However not all affine Hjelmslev planes may be extended to a projective Hjelmslev plane[7]. As a corollary of Theorems 1 and 3 we show that (t, t) -AH planes are well behaved in this respect.

Corrolary 1. *All (t, t) -AH planes a sub-geometry of a (t, t) -PH plane.*

Proof. All orthogonal arrays $OA(2, m, m)$ are extendable to $OA(2, m + 1, m)$ [3, §III Thm 3.95], and hence all (t, t) -AH planes are extendable to $(t, t,)$ -PH planes. $\square$

6 Conclusions

For orders where there are several possible projective planes, affine planes and orthogonal arrays, these algorithms generate many different Hjelmslev planes of the same size. Even with the same seeds, there is an enormous number of choices to be made in step 3, resulting an explosion of the number of planes generated. The problem of determining if two planes are isomorphic is computational intensive. When there is an algebraic structure on the plane, the automorphism group can be very large [14]. Thus it is expected, that a large number of isomorphic planes generated. Further analysis of the choices made in step 3 may reduce the number of isomorphic planes generated. There has been very little research on isomorphisms of Hjelmselv planes, so this is an important direction for further research.

This algorithm has been implement in Python by Jesse Waechter-Cornwill with a visual interpretation.

<http://user.gs.rmit.edu.au/asha/html/resproj.html>.

There is also scope for extending this algorithm to affine planes of higher uniformity by using the 2– uniform affine Hjelmselv plane generated in Algorithm 2, and the corresponding array structure as inputs in Algorithm 1.

Acknowledgement

Thanks to Jesse Waechter-Cornwill for the coding and visualisation of the algorithm 1. Thanks are due to Cathy Baker for highlighting reference [13].

References

- [1] R. A. Bailey, Peter J. Cameron, Peter Dobcsnyi, John P. Morgan, and Leonard H. Soicher. *DesignTheory.org*. U.K. Engineering and Physical Sciences Research Council, designtheory.org, updated on 2009-09-17.
- [2] T. Beth, D. Jungnickel, and H. Lenz. *Design Theory*. Cambridge University Press, 1993.
- [3] C. J. Colbourn and J.H. Dinitz, editors. *Handbook of Combinatorial Designs*. Chapman and Hall/CRC Press, second edition, 2007.
- [4] Robert T. Craig. Extensions of finite projective planes I. Uniform Hjelmslev planes. *Canad. J. Math.*, 16:261–266, 1964.
- [5] P. Dembowski. *Finite Geometries*. Classics in Mathematics. Springer, reprint of the 1968 edition, 1997.
- [6] David A. Drake. On n -uniform Hjelmslev planes. *J. Combin. Theory*, 9:267–288, 1970.
- [7] David A. Drake. Existence of parallelisms and projective extensions for strongly n -uniform near affine Hjelmslev planes. *Geom. Dedicata*, 8:191–214, 1974.
- [8] David A. Drake. Nonexistence results for finite Hjelmslev planes. *Abh. Math. Semin. Univ. Hambg.*, 40:100–110, 1974.
- [9] David A. Drake and Ernest E. Shult. Construction of Hjelmslev planes from $\{t, k\}$ nets. *Geom. Dedicata*, 5:377–392, 1976.

- [10] T. Honold and M. Kiermaier. The existence of maximal $(q^2, 2)$ -arcs in projective hjelmselv planes over chain rings of length 2 and odd prime characteristic. *Designs, Codes and Cryptography*, DOI 10.1007/s10623-012-9653-y, 2012.
- [11] T. Honold and I. Landjev. Linear Codes over Finite Chain Rings. *The Electronic Journal of Combinatorics*, 7(1):R11, 2000.
- [12] Thomas Honold and Ivan Landjev. On arcs in projective Hjelmslev planes. *Discrete Math.*, 231:265–278, 2001.
- [13] G. Hanssens and H₂ Van Maldeghem A universal construction for projective Hjelmselv planes of level n . *Compositio Mathematica*, 71(3): 285-294, 1989.
- [14] M. Koch M. Kiermaier and S. Kurz. 2 arcs of maximal size in the affine and the projective hjelmselv plane over f_{25} . *Advances in Mathematics of Communications*, 5(2):287–301, 2011.
- [15] Erwin Kleinfeld. Finite Hjelmslev planes. *Illinois J. Math.*, 3:403–407, 1959.
- [16] Wilhelm Klingenberg. Projektive und affine ebenen mit nachbarelementen. *Math. Z.*, 60:384, 1954.
- [17] Metod Saniga and Michel Planat. Hjelmslev geometry of mutually unbiased bases. *J. Phys. A*, 39(2):435, 2006.
- [18] N. J. A. Sloane. *A Library of Orthogonal Arrays*. <http://www2.research.att.com/~njas/oadir/>, 2010.
- [19] F.D. Veldkamp. *Handbook of Incidence Geometry*, chapter Geometry over Rings, pages 1033– 1084. Elsevier Science, 1995.